

REVIEW

Open Access

Hyperbaric oxygen treatment for inflammatory bowel disease: a systematic review and analysis

Daniel A Rossignol

Abstract

Background: Traditionally, hyperbaric oxygen treatment (HBOT) has been used to treat a limited repertoire of disease, including decompression sickness and healing of problem wounds. However, some investigators have used HBOT to treat inflammatory bowel disease (IBD), including Crohn's disease and ulcerative colitis.

Methods: Comprehensive searches were conducted in 8 scientific databases through 2011 to identify publications using HBOT in IBD. Human studies and animal models were collated separately.

Results: Thirteen studies of HBOT in Crohn's disease and 6 studies in ulcerative colitis were identified. In all studies, participants had severe disease refractory to standard medical treatments, including corticosteroids, immunomodulators and anti-inflammatory medications. In patients with Crohn's disease, 31/40 (78%) had clinical improvements with HBOT, while all 39 patients with ulcerative colitis improved. One study in Crohn's disease reported a significant decrease in proinflammatory cytokines (IL-1, IL-6 and TNF-alpha) and one study in ulcerative colitis reported a decrease in IL-6 with HBOT. Adverse events were minimal. Twelve publications reported using HBOT in animal models of experimentally-induced IBD, including several studies reporting decreased markers of inflammation or immune dysregulation, including TNF-alpha (3 studies), IL-1beta (2 studies), neopterin (1 study) and myeloperoxidase activity (5 studies). HBOT also decreased oxidative stress markers including malondialdehyde (3 studies) and plasma carbonyl content (2 studies), except for one study that reported increased plasma carbonyl content. Several studies reported HBOT lowered nitric oxide (3 studies) and nitric oxide synthase (3 studies) and one study reported a decrease in prostaglandin E₂ levels. Four animal studies reported decreased edema or colonic tissue weight with HBOT, and 8 studies reported microscopic improvements on histopathological examination. Although most publications reported improvements with HBOT, some studies suffered from limitations, including possible publication and referral biases, the lack of a control group, the retrospective nature and a small number of participants.

Conclusions: HBOT lowered markers of inflammation and oxidative stress and ameliorated IBD in both human and animal studies. Most treated patients were refractory to standard medical treatments. Additional studies are warranted to investigate the effects of HBOT on biomarkers of oxidative stress and inflammation as well as clinical outcomes in individuals with IBD.

Keywords: Hyperbaric oxygen treatment, Inflammation, Oxidative stress, Inflammatory bowel disease, Crohn's disease, Ulcerative colitis, Biomarkers

Background

Inflammatory bowel disease (IBD) is a chronic inflammatory disease of the gastrointestinal (GI) tract characterized by chronic and recurrent ulcerations [1], and includes Crohn's disease and ulcerative colitis. IBD is usually accompanied by severe GI symptoms such as

diarrhea, bleeding, abdominal pain, weight loss, and anemia. The symptoms of IBD can be intermittent, with periods of exacerbations and periods that may be relatively free of symptoms. Recent evidence suggests that the pathophysiology of IBD involves immune dysregulation, genetic susceptibilities, intestinal barrier dysfunction, and alterations in microbial flora [2]. Activated macrophages appear to play a key role in the disease process and produce proinflammatory cytokines,

Correspondence: rossignolmd@gmail.com
Rossignol Medical Center, 3800 West Eau Gallie Blvd., Melbourne, FL 32934, USA